

The liquid formulation of maleic hydrazide for potatoes.

APPLICATION ADVICE



Crop requirements

- Crops should be **actively growing** and free from pest, disease and climatic stress.
 - In late planted crops, consider the maturity of the crop, application is best made 2-3 weeks after flowering.
- Apply from 3 to 7 weeks before haulm desiccation.
 - Some lower leaves may be showing signs of dying off but the canopy should be actively growing.
 - Optimum timing is around **5 weeks before haulm desiccation** to allow even distribution of maleic hydrazide between tubers.
 - Do not apply within 3 weeks of haulm desiccation.
- Apply when sufficient tubers of **size greater than 25 mm** are present to give a marketable yield.
 - Only tubers larger than 25 mm will continue to grow and result in a marketable yield.
- Ensure the **crop foliage is dry**.
- Do not treat potatoes intended for use as seed. Do not treat first early potatoes. Do not treat potatoes grown under polythene.
- Treatments are best made as a **solo application** to maximise plant uptake and translocation to tubers.

Weather conditions

- Apply when conditions are **below 25°C** with a relative humidity of **greater than 50%**.
 - High temperatures can cause maleic hydrazide to crystallise on the leaf surface thereby reducing absorption.
 - High relative humidity greatly increases absorption rate.
- Avoid rainfall and irrigation for **24h** following application.
 - Maleic hydrazide uptake into the leaf is relatively slow, therefore avoid unnecessary wash-off.

Crop	Max individual Dose (L/ha)	Active ingredient (g/ha)	Water volume (L/ha)	Latest timing (days before harvest)
Potato	11.1	2,997	300-600	21
Bulb Onion	8.9	2,403	500-600	4

Itcan SL 270 is available in 15L bottles, 600L and 1,000L IBC's



Kreglinger

Approval holder:
Grote Markt 7
B. 2000 Anvers
Belgium
Tel: 0032 (0)32 222 020
www.kreglinger.com



AGRICULTURE
Marketing company:
71-75 Shelton Street
Covent Garden
London, WC2H 9JQ
Tel: 020 3011 5515
www.geminiag.com

PRODUCTS FOR PROFESSIONALS: USE PLANT PROTECTION PRODUCTS WITH CAUTION. BEFORE USE, READ THE LABEL AND INFORMATION RELATING TO THE PRODUCT.

Itcan SL 270® - SL, Soluble Concentrate - 270 g/l of maleic hydrazide – Follow the instructions for use in order to avoid risks to human health and the environment - Do not pollute water with the product or its packaging - Registered trademark: Drexel - Authorisation holder: Kreglinger Europe N.V. - Read the label carefully before any use, and comply strictly with usages, doses, conditions and precautions for use.



@geminiagriculture



@gemini_ag

